



Opportunities and challenges in using remaining carbon budgets to guide climate policy

H. Damon Matthews¹✉, Katarzyna B. Tokarska², Zebedee R. J. Nicholls^{3,4}, Joeri Rogelj^{5,6}, Josep G. Canadell⁷, Pierre Friedlingstein^{8,9}, Thomas L. Frölicher^{10,11}, Piers M. Forster¹², Nathan P. Gillett¹³, Tatiana Ilyina¹⁴, Robert B. Jackson^{15,16}, Chris D. Jones¹⁷, Charles Koven¹⁸, Reto Knutti¹⁹, Andrew H. MacDougall¹⁹, Malte Meinshausen¹⁹, Nadine Mengis^{20,21}, Roland Séférian²² and Kirsten Zickfeld²¹

The remaining carbon budget represents the total amount of CO₂ that can still be emitted in the future while limiting global warming to a given temperature target. Remaining carbon budget estimates range widely, however, and this uncertainty can be used to either trivialize the most ambitious mitigation targets by characterizing them as impossible, or to argue that there is ample time to allow for a gradual transition to a low-carbon economy. Neither of these extremes is consistent with our best understanding of the policy implications of remaining carbon budgets. Understanding the scientific and socio-economic uncertainties affecting the size of the remaining carbon budgets, as well as the methodological choices and assumptions that underlie their calculation, is essential before applying them as a policy tool. Here we provide recommendations on how to calculate remaining carbon budgets in a traceable and transparent way, and discuss their uncertainties and implications for both international and national climate policies.

Remaining carbon budgets are defined as the allowable future CO₂ emissions that are consistent with meeting climate targets such as those of the Paris Agreement (see Box 1). Conceptually, the idea of a global emissions budget is a compelling way to frame and communicate the climate mitigation challenge: a finite cap on total CO₂ emissions implies clearly that global CO₂ emissions must eventually reach net-zero to stabilize global temperatures. Estimates of the remaining carbon budget are subject to large uncertainty, but have also varied considerably among studies owing to the lack of a consistently applied definition, as well as different methodological approaches used to calculate the remaining budgets^{1,2}. Furthermore, additional uncertainties are introduced in the process of disaggregating the global budget into national shares for domestic climate policy^{3–5}. Given the increasing adoption of remaining carbon budget estimates as a benchmark for national policy discussions, the full range of uncertainties and choices surrounding carbon budget estimates must be articulated and understood.

In this Perspective, we present an overview of the state of our understanding of the remaining carbon budget, with the intent of charting a tractable path through the scientific, policy and ethical

considerations required when applying the carbon budget concept to climate policy decisions. We characterize the uncertainties and other factors affecting estimates of the remaining carbon budget across four broad categories: (1) geophysical uncertainties associated with physical climate and carbon cycle processes that determine the climate response to emissions; (2) socio-economic uncertainties that reflect the societal choices and dynamics that determine future emission scenarios; (3) methodological approaches that reflect choices and assumptions made when estimating the remaining carbon budget; and (4) allocation choices that emerge from the range of ethical and fairness principles that can be used to allocate a portion of the global budget to individual countries, economic sectors and entities such as individual industries and corporations. We discuss each of these in turn, and then offer some concluding thoughts on the robust policy implications of a finite remaining carbon budget.

Geophysical basis of carbon budgets

The proportionality between cumulative CO₂ emissions and CO₂-induced temperature change is the primary geophysical basis

¹Department of Geography, Planning and Environment, Concordia University, Montréal, Canada. ²Institute for Atmospheric and Climate Science, ETH Zürich, Zürich, Switzerland. ³Australian–German Climate and Energy College, The University of Melbourne, Melbourne, Australia. ⁴School of Earth Sciences, The University of Melbourne, Melbourne, Victoria, Australia. ⁵Grantham Institute – Climate Change and Environment, Imperial College London, London, UK. ⁶International Institute for Applied Systems Analysis (IIASA), Laxenburg, Austria. ⁷Global Carbon Project, CSIRO Oceans and Atmosphere, Canberra, Australian Capital Territory, Australia. ⁸College of Engineering, Mathematics and Physical Sciences, University of Exeter, Exeter, UK. ⁹Laboratoire de Météorologie Dynamique, Département de géosciences, École Normale Supérieure, PSL Research University, Sorbonne Université, École Polytechnique, CNRS, Paris, France. ¹⁰Climate and Environmental Physics, Physics Institute, University of Bern, Bern, Switzerland. ¹¹Oeschger Centre for Climate Change Research, University of Bern, Bern, Switzerland. ¹²Priestley International Centre for Climate, University of Leeds, Leeds, UK. ¹³Canadian Centre for Climate Modelling and Analysis, Environment and Climate Change Canada, Victoria, British Columbia, Canada. ¹⁴Max Planck Institute for Meteorology, Hamburg, Germany. ¹⁵Department of Earth System Science, Woods Institute for the Environment, Stanford University, Stanford, CA, USA. ¹⁶Percourt Institute for Energy, Stanford University, Stanford, CA, USA. ¹⁷Met Office Hadley Centre, Exeter, UK. ¹⁸Climate and Ecosystem Sciences Division, Lawrence Berkeley National Lab, Berkeley, CA, USA. ¹⁹St. Francis Xavier University, Antigonish, Nova Scotia, Canada. ²⁰GEOMAR, Helmholtz Centre for Ocean Research, Kiel, Germany. ²¹Simon Fraser University, Burnaby, British Columbia, Canada. ²²CNRM, Université de Toulouse, Météo-France, CNRS, Toulouse, France.

✉e-mail: damon.matthews@concordia.ca

Box 1 | What is a 'remaining carbon budget'?

Here, a **carbon budget** is the quantity of cumulative or total CO₂ emissions that is consistent with limiting global mean warming to a given temperature level. The term 'budget' is analogous to a fixed total financial budget in that spending in excess of annual budget amounts in the near-term requires decreased spending in the future to not exceed the total budget. Here, we distinguish between the total and the remaining carbon budget: (1) the **total carbon budget** is defined as the total amount of CO₂ emissions since the pre-industrial reference period that is consistent with a specified peak temperature increase; and (2) the **remaining carbon budget** represents the amount of CO₂ that can still be emitted from the present day onwards while staying below the temperature target^{1,2}. In both cases, these budgets refer to a total quantity of CO₂ emissions (from fossil fuels and land-use change) up to the point in time that CO₂ emissions reach net-zero. Importantly, this quantity applies to only CO₂ emissions, and does not apply to allowable CO₂-equivalent emissions of other gases and aerosols (for example, methane and nitrous oxide, as calculated using global warming potentials).

There are several variants of carbon budgets that have been used in the literature (for example, **threshold exceedance budget**,

threshold avoidance budget, **threshold return budget** and **overshoot budget**; see refs. ^{1,38} for a discussion of these variants). We do not use these terms further, and we recommend that they be used only if necessary to explore the sensitivity of carbon budget estimates to different types of emission scenarios or estimation methods.

Another common but distinct use of the term 'carbon budget' is to describe the balance of historical sources and sinks of CO₂ in the Earth system or for a particular region. In this alternative usage, the term 'budget' is similar to annual financial accounting in a closed system without deficits, whereby the sum of the individual positive and negative line items must equal zero. As a prominent example, the Global Carbon Project annually publishes a **historical global carbon budget**, in which they assess the human perturbation to the global carbon cycle up to the present day⁹¹. This historical global carbon budget combines fossil fuel and land-use CO₂ emissions (or sources) with increases in atmospheric, land and ocean carbon reservoirs (also referred to as sinks) and places these sources and sinks in the context of natural carbon cycle processes.

for a finite remaining carbon budget: each additional tonne of CO₂ emitted leads to an incremental temperature increase, which implies that CO₂ emissions must decrease to net-zero to stabilize global temperature⁶. This proportionality is quantified by the transient climate response to cumulative CO₂ emissions (TCRE), which defines the transient warming per unit cumulative CO₂ emissions in a scenario with increasing CO₂ emissions (see Box 2). The allowable cumulative CO₂ emissions for a given amount of warming are therefore proportional to the inverse of the TCRE^{1,7}. However, this relationship only holds for the warming caused by CO₂, and not for additional warming (or cooling) caused by other emissions or forcings (for example, methane, aerosols or nitrous oxide).

Using the TCRE to estimate the remaining carbon budget for a given temperature target therefore requires an additional estimate of the non-CO₂ contribution to future warming^{1,7–9}. One approach is to define an 'effective TCRE' to estimate the total anthropogenic warming at a given amount of cumulative CO₂ emissions^{9–11} (see Box 2). However, unlike the TCRE, there is no geophysical basis for the effective TCRE to remain constant in time. In particular, where the TCRE is approximately scenario-independent^{12,13}, this does not hold for the effective TCRE, which is affected by the changing rate of emissions of non-CO₂ forcings with mostly shorter atmospheric lifetimes than that of CO₂¹⁴. Consequently, while the effective TCRE can be used to calculate the total carbon budget directly¹⁰, it is important to use an estimated value of the effective TCRE at the time that the temperature target is reached in a given scenario (Fig. 1).

Geophysical uncertainties

Geophysical uncertainties affecting the TCRE (see Box 2) will propagate directly to uncertainties in estimates of the remaining carbon budget¹⁵. In general, there are three types of geophysical uncertainties relevant to the TCRE that affect its robustness as a predictor of the remaining carbon budget. First, differing physical climate and carbon cycle responses to CO₂ emissions alter the magnitude of the TCRE. Across Earth-system models, TCRE values are generally proportional to a model's transient climate response (TCR), though different carbon cycle responses will also produce a range of TCRE values among models with similar TCR values^{16,17} (see Box 2). Better constraints on the TCR value (see, for example, refs. ^{18,19}) as well as on the carbon cycle response to climate change and increasing CO₂

would therefore have the effect of constraining both the TCRE and the remaining carbon budget.

Second, the linearity of the TCRE relationship results from the compensation of individual non-linear processes that act to both increase and decrease the sensitivity of the temperature response to additional cumulative CO₂ emissions^{20–23}. However, there is nevertheless the potential for strong non-linear changes in the strength of particular feedbacks to cause deviations from the TCRE-predicted linearity with increasing emissions. Examples include potential changes to the strength of physical climate feedbacks as a result of changing warming patterns^{24–26} or the behaviour of biogeochemical permafrost and wetland feedbacks that are currently poorly represented in Earth-system models^{27,28}.

Third, while the TCRE has been shown to be a robust predictor of CO₂-induced warming in scenarios with increasing CO₂ emissions^{29,30}, it is less clear that the TCRE will adequately capture the climate response to scenarios that rapidly decline to net-zero and/or net-negative CO₂ emissions^{31,32}. In the event that there is unrealized warming or cooling from past CO₂ emissions, this lagged temperature change would manifest during the time that CO₂ emissions ramp down to zero, causing the temperature response to cumulative emissions to bend upwards or downwards relative to the linear TCRE line. This unrealized commitment from past CO₂ emissions has recently been quantified across Earth-system models³³, and is probably an important contributor to uncertainty in the remaining carbon budget¹⁵.

Finally, the climate response to non-CO₂ forcing changes is an important additional source of geophysical uncertainty affecting carbon budget estimates that is distinct from the contributions to TCRE uncertainty discussed above (see 'Non-CO₂ forcing and climate response' bar below Fig. 1a; ref. ⁷). Given the prominence of aerosol forcing uncertainty in affecting the overall non-CO₂ forcing uncertainty, this speaks to the importance of improved estimates of the climate response to present-day aerosol forcing in order to improve our ability to constrain estimates of the remaining carbon budget.

Socio-economic scenario uncertainty

Remaining carbon budget estimates are also strongly affected by socio-economic uncertainties related to our ability to predict the dynamics of socio-political systems and the technological changes


Box 2 | The transient climate response to cumulative CO₂ emissions (TCRE)

A close to proportional relationship between CO₂-induced global warming and cumulative CO₂ emissions is an emergent property of a range of Earth-system models^{12,13,77,92,93}. The TCRE quantifies the temperature change per unit of cumulative CO₂ emissions (often expressed as °C per 1,000 GtC or per 1,000 GtCO₂ emitted). In a climate model, the TCRE defines the temperature change per unit CO₂ emissions at the time of doubled atmospheric CO₂ concentration, in an idealized experiment where atmospheric CO₂ concentration increases at a rate of 1% per year^{29,30}. This TCRE value can then be used to describe the general linear relationship between cumulative CO₂ emissions and CO₂-induced temperature change (Fig. 1a, solid line, where the value of the TCRE defines the slope of this line; see also Box 2 figure, panel a).

The TCRE has been shown to be a good predictor of warming caused by a given quantity of cumulative CO₂ emissions across a range of emissions scenarios¹³, though it is subject to many uncertain processes that affect both its magnitude (the slope of the line) and its constancy in time (the robustness of the linear response to additional cumulative emissions) (Box 2 figure, panel a). Among Earth-system models, the magnitude of the TCRE is strongly related to the TCR, which accounts for at least half of the variation in TCRE values^{16,17}; the remaining TCRE variation is caused by varying carbon cycle sensitivity to global warming and increasing CO₂¹⁷ (Box 2 figure, panel b). Model TCRE values are also (though to a lesser extent) related to their equilibrium climate sensitivity (ECS; the long-term global surface warming that is expected to occur in response to a doubling of the atmospheric CO₂ concentration). Several models in the recent Sixth Coupled Model Intercomparison Project⁹⁴ (CMIP6) have higher ECS values than the previous generation of models⁹⁵ (CMIP5), most of which are also associated with high TCR and TCRE values. However, some

of these high-ECS CMIP6 models have TCR and TCRE values that fall outside of estimates of the observationally constrained 5–95% range^{18,19,29} (Box 2 figure, panel b). Consequently, while these high-ECS models would predict smaller remaining carbon budgets, this should be considered a low-probability outcome given their current lower consistency with observed warming^{18,96}. Such results can nevertheless be used to guide quantitative risk assessment of the implications of our imperfect knowledge of climate processes and the associated risks of low-probability outcomes⁹⁷.

Importantly, the TCRE applies only to warming caused by CO₂ emissions, and does not include the additional warming or cooling caused by non-CO₂ emissions and other climate drivers. A common approach to incorporate the effect of non-CO₂ forcing is to use simulations forced by all anthropogenic drivers, and plot the resulting total anthropogenic warming as a function of cumulative CO₂ emissions. This results in a representation of total anthropogenic warming per unit cumulative CO₂ emission that includes the effect of all climate drivers^{45,98}. Here, we label this the effective TCRE (following ref. ¹⁰) to define the total anthropogenic warming at a given quantity of cumulative CO₂ emissions (Box 2 figure, panel a, slope of dashed black line). The effective TCRE is a measure of warming per unit cumulative CO₂ emissions, equal to the TCRE plus an additional amount of scenario-dependent non-CO₂ warming. This metric has been used conceptually or explicitly in several previous studies to directly infer remaining carbon budgets associated with a given future emission scenario^{9–11,45,81–83}. However, this needs to be done with an understanding that the effective TCRE is not expected to remain constant in time (see Fig. 1) due to its strong dependence on non-CO₂ scenario variation and in particular on the potential unmasking of aerosol cooling in scenarios with rapid decreases in aerosol emissions¹⁴.



Uncertain processes affecting the relationship between warming and cumulative CO₂ emissions. **a**, Conceptual representation of the TCRE (the slope of the solid black line) and the effective TCRE (represented by the dashed black line). Solid coloured arrows indicate scientific uncertainties affecting the value of the TCRE due to the climate (pink) and carbon cycle (orange) response to CO₂ emissions^{16,29}. Solid coloured lines indicate processes that may lead to deviations from a linear temperature response to cumulative emissions. Notable processes are the additional unrealized warming (or cooling) due to past emissions that could occur as CO₂ emissions approach zero^{33,99} (CO₂ commitment; purple), permafrost carbon feedbacks^{27,31} (light green), and the increasing saturation of CO₂ radiative forcing at high CO₂ levels²⁰ (dark green). Other nonlinear climate or carbon cycle processes act in both directions and tend to compensate for one another (turquoise), leading to an approximately constant TCRE value across a wide range of cumulative CO₂ emissions^{12,13,21,29,84}. Dashed coloured lines indicate additional uncertainties affecting the effective TCRE as a result of societal choices leading to different non-CO₂ greenhouse gas emission scenarios^{7–9,14,45} (dashed blue) and the warming response to decreased aerosol emissions^{34,35} (dashed red). **b**, Relationship between model TCRE and TCR values in the CMIP5 models^{29,100,101} (triangles) and CMIP6 models^{16,18} (circles). Grey shading indicates the observationally constrained TCR median (dashed line), probable range (>66%; dark grey) and 5–95% range (light grey), based on the 1981–2017 observed warming trend as a constraint based on ref. ¹⁸. The blue bar and blue dotted lines indicate the observationally constrained TCRE range (5–95%) reported by ref. ²⁹.




Fig. 1 | Relationship between the TCRE, the effective TCRE, and the total and remaining carbon budgets. a, Idealized representation of the TCRE, effective TCRE and related total and remaining carbon budgets. Here, we show the central estimate of the TCRE (0.45 °C per 1,000 GtCO₂) and an effective TCRE value of 0.53 °C per 1,000 GtCO₂ at 1.5 °C, which corresponds to the median (50th percentile) remaining carbon budget of 580 GtCO₂ from 2018 onwards reported in the IPCC Special Report on Global Warming of 1.5 °C (SR1.5; table 2.2. of ref. ⁷). The 67th (420 GtCO₂) and 33rd (840 GtCO₂) percentile budgets are shown in lighter red, reflecting uncertainty in the TCRE only. Other contributions to uncertainty are shown by bars below the plot, showing the amounts by which these additional processes affect the median budget estimate (table 2.2. of ref. ⁷). **b**, Simulated climate response to CO₂ emissions only (purple) and all anthropogenic drivers (red) for scenarios from the SR1.5 scenario database^{87,88}, using the simple model emulator MAGICC7 (refs. ^{89,90}) with parameter settings corresponding to a TCRE of 0.44 °C per GtCO₂. Temperature change is shown relative to the 1850–1900 period, and cumulative CO₂ emissions are calculated from the central year of that period (1875). Dots mark the peak warming and the lines end at the point of net-zero CO₂ emissions in each scenario. The larger spread of red dots relative to purple shows the additional effect of socio-economic uncertainty on the effective TCRE as a result of differing non-CO₂ emission scenarios.

that determine the evolution of emissions in future scenarios¹⁵. Given that the TCRE is relatively robust to variation in CO₂ emission scenarios^{13,20,23}, the relationship between the TCRE and the remaining carbon budget is only affected by socio-economic uncertainty in the case of a large positive or negative zero emissions commitment (ZEC). However, the effect of socio-economic uncertainty is considerably more pronounced for estimates of future non-CO₂ warming (see spread of total warming as a function of cumulative emissions across scenarios in Fig. 1b (red dots)). This relates to the shorter atmospheric lifetimes of many non-CO₂ emissions, such that rapid decreases in emissions of short-lived positive climate forcings (such as methane, black carbon and ozone precursors) would effectively limit near-term warming caused by non-CO₂ emissions¹⁴. Conversely, however, rapid decreases in aerosol emissions that produce a negative forcing would amplify non-CO₂ warming¹⁴, a scenario that is likely to occur as a result of decarbonization efforts^{34–36}.

Although both geophysical and socio-economic uncertainties can be reduced by further research, socio-economic uncertainties are also sensitive to human decisions and choices regarding technological development and mitigation actions³⁷. This means that

policy decisions about where to focus mitigation efforts have the potential to influence the size of the remaining carbon budget by affecting the amount of warming that is caused by CO₂ versus other anthropogenic climate drivers^{37,38}. The balance of effective mitigation of positive short-lived forcings and the potential aerosol warming commitment may be one of the most important determinants of the size of the remaining carbon budget.

Methodological choices and assumptions

Using a consistent and transparent set of assumptions to calculate the remaining carbon budget is crucial in order to provide clear guidelines for climate policy. Yet, inconsistent choices are often used among different studies that report remaining carbon budget estimates, which unnecessarily inflates the spread of estimates^{1,2}. Here, we provide guidelines and recommendations to estimate carbon budgets in a transparent and policy-relevant way, given the many choices and assumptions typically required (Table 1).

First, we recommend estimating remaining carbon budgets for anthropogenic warming only, independent of natural variability (for example, as estimated in ref. ³⁹); while this may seem like an obvious

Table 1 | Choices and assumptions that are typically required when estimating total or remaining carbon budgets for global temperature targets

Choice	Options	Issues to consider
Definition of global warming	(*) Anthropogenic warming only	Relevant to international climate targets aimed at limiting global temperature increase ⁴⁰ ; requires a method to isolate the anthropogenic contribution to observed or model-simulated temperature change ^{29,39} .
	Anthropogenic warming + natural climate variability	Directly observable and simulated by global climate models; however, natural climate variability, whether externally forced (by volcanoes or solar activity) or unforced (that is, internal variability of the climate system), causes inter-annual to decadal-scale warming or cooling trends that are not relevant to international climate goals ^{40,42,43} .
Target temperature	(*) 1.5 °C	Most ambitious target level in the Paris Agreement.
	(*) "well below 2 °C"	Primary Paris Agreement target, but not precisely defined.
	2 °C or higher	Warming levels that exceed the Paris Agreement target range.
Probability of not exceeding the target	50%, 67%, 90%, ...	Remaining carbon budgets are typically estimated to be in line with a 50%, 67% or sometimes 90% probability of successfully limiting warming to the temperature threshold of interest. The choice of which budget to adopt as a global target depends on societal risk-avoidance preferences.
Pre-industrial reference period	(*) 1850–1900 average	Current proxy for pre-industrial climate ⁴⁶ , corresponding with the beginning of available instrumental temperature records.
	1860–1880 average	Period representing the first 20 years available in the HadCRUT temperature dataset ⁷² , at times used because no major volcanic eruption took place during these years.
	1720–1800 average	Suggested by ref. ⁴⁷ as a better estimate of climate conditions prior to human influence, but direct observations of global temperature are not available and emissions uncertainty prior to 1850 is very large, posing difficulty for consistent estimates of historical cumulative emissions from earlier time periods.
Temperature change metric	Observed (blended air–water and/or masked) temperature (GMST)	Directly observable; calculated as a combination of surface air temperatures over land and sea ice, with surface water temperature over ocean; incomplete spatial coverage ⁷³ unless infilling technique used to extrapolate to areas with no observations ⁷⁴ ; spatial definition changes over time as sea ice cover changes ² .
	GSAT	Average of surface air temperature with complete global coverage; typical output of global models, but not currently available from observational estimates of global temperature change; historical GSAT warming has been estimated to be about 0.1 °C higher than that based on GMST ^{75,76} , though recent improvements to spatial infilling techniques in GMST products have decreased this difference.
Nature of temperature change	(*) Peak temperature	The timing of peak temperature should match closely the timing of net-zero CO ₂ emissions ⁷ , which avoids the need for assumptions related to the reversibility of temperature overshoots.
	Temperature at some level of cumulative emissions	No strict limit on maximum temperature change: temperature may exceed the target after the point at which cumulative emissions are calculated.
	Temperature at some year (for example, 2100)	No strict limit on maximum temperature change: temperature may exceed target either before (overshoot and return scenarios) or after selected year.
Non-CO₂ scenario choice	(*) Non-CO ₂ scenario consistent with CO ₂ emissions that decline to net-zero	Requires an internally consistent CO ₂ and non-CO ₂ emission scenario that is also consistent with the desired target, and/or an embedded economic model that generates consistent CO ₂ and non-CO ₂ emissions ⁷⁶⁰ .
	Non-CO ₂ forcing consistent with high-emissions (or other arbitrary) emission scenario	Non-CO ₂ forcing is not consistent with mitigation efforts to limit warming to the target, potentially leading to either an under- or overestimate of the remaining carbon budget ^{8,9,31} .
Treatment of aerosol forcing uncertainty	Single model run for time-series of historical and future aerosol forcing	Results are contingent on the model and/or forcing scenario used and do not include uncertainty associated with aerosol forcing or the climate response to decreased future aerosol emissions ⁵ .
	Assumed range of forcing	Requires many model simulations or other methods to sample uncertainty ⁷⁷ .
Treatment of land-use CO₂ emissions	Prescribed or treated as interchangeable with fossil fuel emissions	Cumulative emissions from land use are known, but biophysical and indirect carbon cycle effects of land-use change are not accounted for.
	Simulated by model from prescribed land-use change scenario	Biophysical and other effects of land-use change accounted for, but cumulative emissions from land use are complex to diagnose, often requiring additional simulations ^{78–80} . In the absence of diagnosed land-use emission information, some studies have assumed the same land-use change emissions in all model simulations ^{81–83} .

Continued

Table 1 | Choices and assumptions that are typically required when estimating total or remaining carbon budgets for global temperature targets (continued)

Choice	Options	Issues to consider
Treatment of CO₂ ZEC	Assumed to be zero or negligible	This is the approach taken in ref. ⁷ , but results in unquantified uncertainty associated with the ZEC.
	Assumed to have a positive or negative value	When combined with a method that uses TCRE directly, a positive ZEC would reduce the remaining carbon budget to account for unrealized warming after CO ₂ emissions are halted ^{1,84} . Similarly, a negative ZEC would increase the remaining budget, but only if its timescale is shorter than the pace of emission reductions to net-zero CO ₂ .
	Simulated by model using scenarios where CO ₂ emissions decrease to net-zero	The effect of the CO ₂ ZEC is accounted for in simulations of ambitious mitigation scenarios, as any unrealized warming or cooling from past emissions would typically be realized during the time that emissions fall to zero.
Treatment of overshoot scenarios/ reversibility	Climate response to CO ₂ removal assumed to be the same as response to emissions	Many recent studies are based on this assumption (see ref. ¹ for a summary of different studies). This assumption is not supported if the CO ₂ warming commitment (ZEC) is non-zero ⁶¹ .
	Overshoot scenario simulated explicitly	Resulting budget is specific to overshoot scenarios ^{31,32} .
	Overshoot scenarios excluded	Resulting budget may not apply to overshoot scenarios ²⁷ .
Treatment of permafrost feedback	Included in model simulation, or using TCRE from models that include this feedback	Resulting budget does not need to be adjusted for this feedback ³¹ , but timescale is important as permafrost feedbacks have a long time constant ^{85,86} .
	Not included in model or using TCRE range from CMIP5 models	Budget should be adjusted to account for the effect of this feedback, and timescale needs to be stated given the feedback's time dependency ⁷ .
Treatment of other under- or unrepresented feedbacks	Clarity about which feedbacks are included and which are not	Representation and strength of feedbacks vary among models, which is part of the explanation for the existing TCRE range. In general, any new positive (amplifying) feedback would increase the TCRE (and decrease the budget), and a new negative (attenuating) feedback would decrease the TCRE (and increase the budget) ¹ .

Where appropriate, recommended choices are marked with an asterisk (*).

statement, it is nevertheless the case that climate policy temperature goals have not been consistently interpreted across different studies⁴⁰. The ultimate objective of the United Nations Framework Convention on Climate Change (UNFCCC) is to prevent “dangerous anthropogenic interference in the climate system” (Article 2)⁴¹. This framing provides a clear rationale for limiting warming caused by all anthropogenic drivers, rather than that caused by the combined effect of anthropogenic warming and natural variability^{40,42,43}.

Second, we recommend that remaining carbon budgets be defined in relation to a particular policy-relevant climate target. The Paris Agreement⁴⁴ aims to keep global temperatures to “well below 2 °C above pre-industrial levels” while “pursuing efforts to limit the temperature increase to 1.5 °C above pre-industrial levels”. Consequently, carbon budgets associated with a range of temperature increases between 1.5 °C and “well below 2 °C” are those with direct relevance for the Paris Agreement goal. Budgets for 2 °C or higher can be used to gauge the amount of effort needed to stay below higher warming levels, but are outside the current policy framing of the Paris Agreement.

Third, choices about the desired level of risk avoidance must be defined when estimating remaining carbon budgets. Carbon budgets have typically been reported as corresponding to a 50% or 67% percent probability of staying below a given temperature target when the carbon budget is fully emitted, given known and quantified uncertainties⁴⁵.

Fourth, the proxy for the pre-industrial reference period that has been used in recent Intergovernmental Panel on Climate Change (IPCC) reports^{7,46} is the 1850–1900 average, which we recommend

adopting for consistency with these analyses. We recognize that temperatures during this period may have already increased relative to the previous century^{39,47,48}, though adopting an earlier baseline period is currently difficult owing to limited observational data on both temperature and cumulative CO₂ emissions prior to 1850.

Fifth, it is important to be explicit about the choice of temperature-change metric. The relative merits of using a blended air–water temperature that is masked according to observational coverage (global mean surface temperature, GMST), a full-coverage global surface air temperature (GSAT) or a combination of the two to represent observed historical warming have been discussed extensively elsewhere^{1,2,49}; we do not offer a specific recommendation here other than that the choice of metric be clear and justified, both for historical warming and for the value of the TCRE.

Sixth, we recommend adopting our carbon budget definition of total emissions up to the point in time that CO₂ emissions reach net-zero (see Box 1), which is likely to correspond closely to the timing of peak temperature change⁷. This choice also avoids the need for assumptions regarding the potential feasibility of net negative CO₂ emissions to reverse temperature overshoots. For model simulations, this requires using emissions scenarios that contain internally consistent CO₂ and non-CO₂ emissions, which are also broadly consistent with a desired peak temperature target, rather than scenarios where temperature exceeds the target indefinitely or exceeds (overshoots) and returns to the target in question.

For the other methodological choices listed in Table 1, our key recommendation is for each choice to be documented to clarify the assumptions embedded in analyses, and to discuss (quantitatively if possible) how choices may affect the results. Lack of clarity




Fig. 2 | Illustrative example of distributing the remaining carbon budget over time into five-year discrete time intervals. The years of net-zero CO₂ emissions shown here are approximately consistent with the estimates of the 67th percentile remaining budgets from SR1.5' (420 GtCO₂ after 2017; area under the blue line) and for an illustrative "well below 2 °C" interpretation, here taken to be 1.75 °C (800 GtCO₂ after 2017; area under the yellow line). The red shaded regions indicate projected amounts over budget, reflecting estimates of global CO₂ emissions between now and 2030 following current NDCs⁵⁰. We note that the linearly decreasing trajectories illustrated here are clearly an idealized scenario and do not incorporate aspects of cost-effectiveness; however, the small size of remaining carbon budgets for the Paris Agreement goal clearly requires stronger reductions in global CO₂ emissions in the next decades than are captured by current NDCs⁵⁰.

with respect to these assumptions and choices can result in widely varying carbon budget estimates that risk being applied inappropriately to policy questions. For example, carbon budget estimates that assume non-CO₂ forcing will follow a high-emission scenario should not be applied uncritically to the case of ambitious mitigation scenarios with decreasing non-CO₂ emissions. Similarly, budget estimates using the GMST temperature metric are not well suited to estimating the requirements for avoiding climate impacts that have been calculated using GSAT change. Consistent and clear methodological choices are therefore critical to minimizing the risk of misuse of carbon budget estimates in policy applications.

Application to international and national climate policy

Carbon budgets are a powerful guide for international policy, providing a way to gauge the consistency of national emission targets with global temperature goals. To do so, the total quantity of remaining CO₂ emissions must first be distributed over time so as to align with the target years of national emission pledges. Figure 2 illustrates one such time distribution, in which a scenario of linearly decreasing CO₂ emissions to zero at 2040 (for a 1.5 °C carbon budget) or 2060 (for "well below 2 °C" budget) is discretized into five-year budgets that are compatible with the remaining carbon budget. Such subdivided budgets could be tracked and used to inform the five-year global stocktake process as part of the implementation of the Paris Agreement, whose mandate is to assess the consistency of emissions targets with the long-term temperature goal. Currently, emissions from national pledges are expected to exceed the near-term budget allowances shown in Fig. 2^{50,51}. This raises important questions of intergenerational equity as we are either accruing an emissions debt to future generations by borrowing allowable emissions from the future allowance, tasking future generations with the challenge

of removing anthropogenic CO₂ from the atmosphere, or committing these future generations to climate change in excess of our stated climate target.

To be useful as a benchmark for national climate policy, the global remaining carbon budget must be further subdivided among nations. There are many choices involved in the distribution of the remaining global carbon budget to individual nations; such decisions often reflect different ways of accounting for unequal national circumstances^{3,4,52–55} (see Box 3). Given the contentious and highly context-specific nature of such distributions, the UNFCCC has chosen not to develop rules or instruct nations as to how to set their own emissions targets. National carbon budgets are therefore currently unilateral choices that can help nations to organize their mitigation efforts, but may not bear any real resemblance to or consistency with the overall global budget. Currently, many nations have designed their nationally determined contributions (NDCs) using the most generous of available allocation principles for their country⁵³. This suggests a need for international mechanisms to promote the evaluation and iterative reassessment of national budget allowances to ensure consistency with the global budget (Box 3).

Despite the challenges associated with fairly allocating the global budget to nations, the idea of a finite national carbon budget nevertheless has enormous conceptual importance for national policy decisions. For example, short-term carbon budgets could be adopted and reported on in a manner similar to fiscal budgets⁵⁶; the simple act of doing so has the potential to embed national climate targets in a much more tangible way across government decision-making processes. The recent adoption of national carbon budgets in the United Kingdom and the European Union, including commitments to net-zero emissions by 2050, are good examples of such an approach. The adoption of finite national budgets would also lend new weight to discussions surrounding new infrastructure construction, and particularly new fossil fuel energy infrastructure. Two recent analyses suggest that global 'committed emissions' associated with existing infrastructure (that is, the future emissions that are expected to occur over the typical operating lifetime of existing infrastructure) are close to or exceed the remaining carbon budgets for our most ambitious climate targets^{57,58}. The committed emissions associated with new infrastructure projects could therefore be weighed against a country's remaining carbon budget to determine whether the infrastructure in question is consistent with our climate targets. Furthermore, should a given nation's emissions exceed their share of the global budget, the resulting emission debts^{4,59} could be used as a metric to inform decision-making related to international climate finance.

Policy implications of a finite remaining carbon budget

The most important policy implication of a finite carbon budget is the need to achieve net-zero CO₂ emissions in order to stabilize global temperature. This framing has been used to inform ongoing and ambitious mitigation efforts in 71 countries and over 100 cities, as well as over 500 businesses that have set net-zero emission targets for a specified year⁵⁰. The concept of a carbon budget has therefore been an effective communication tool in mobilizing individual countries or regions to set net-zero emission reduction targets. However, it is important to reiterate that in order to be consistent with Paris temperature goals, a net-zero CO₂ emission target must also be accompanied by aggressive mitigation of non-CO₂ emissions such as methane, nitrous oxide and black carbon. Both net-zero CO₂ emissions and decreasing non-CO₂ forcing are required to achieve stable global temperatures⁶⁰.


Another important policy implication is that the size of the remaining carbon budget is sensitive to societal choices for mitigating non-CO₂ emissions, as well as the effect of CO₂ mitigation efforts on co-emitted non-CO₂ species^{8,9,31,37}. However, while the remaining

Box 3 | Issues of fairness and equity when allocating the remaining carbon budget to countries

Estimates of the remaining carbon budget provide a global envelope within which future societies have to operate if we intend to limit global warming to a specific level. However, translating this **global budget** to **national allocations** that define what would be an appropriate or fair share of this budget for a single country is an exercise fraught with value judgments that have little relation to the geophysical underpinnings of carbon budgets. Here, science can at best inform and quantify the implications of what are largely subjective choices by individual countries.

Over time, many so-called fairness or equity principles have been suggested and explored to try to understand what would be a fair allocation of the remaining carbon budget. These principles are largely based on concepts of responsibility, equality and capability across nations¹⁰² (Box 3 figure). **Responsibility** addresses the fact that countries have contributed differently to the warming we are currently experiencing, and have also had access to varying levels of understanding over time about the impact of greenhouse gas emissions on the climate. **Equality** reflects the idea of the human right to development, and that each individual is entitled to equal access to the means of development. This principle can be used to imply an equal entitlement to the production of greenhouse gas emissions, though it is also the case that development can be achieved via low-emission technologies and activities. **Capability** reflects the fact that different countries can be in quite different positions regarding their capacity to address the challenge of climate change mitigation, be it in terms of financial resources, technical expertise or institutional context. Historically, this capability has been closely related to a country's degree of industrialization, and its associated greenhouse gas emissions. These various principles are well established in the UNFCCC, which states that countries should participate in responding to climate change “in accordance with their common but differentiated responsibilities and respective capabilities and their social and economic conditions”⁴¹.

A wide variety of ways exist to translate these equity principles by means of quantitative proxies into a distribution method to allocate the global remaining carbon budget to particular countries⁵³. The resulting allocations can be positive or negative; that is, they can represent either an emissions allowance or an emissions debt. Allocation methods also vary considerably in



Allocating the global remaining carbon budget to individual nations. Any such allocation requires subjective choices and application of fairness principles related to a country's responsibility for climate changes, its capability to achieve mitigation goals, and the importance of equality among countries. Coherence between the sum of national allocations and the global allowable budget is unlikely to emerge from this allocation process, though could be achieved with additional evaluation and iterative reassessment of national allocations.

their degree of fairness, and some are generally considered to be explicitly unfair. For example, an equal-shares (sometimes called grandfathering) allocation approach is often used to claim future emissions rights based on the current distribution of emissions among countries. This allocation method does not account for differing historical responsibility or the history of colonial relationships among countries, and instead rewards historical polluters for their current high share of global emissions.

Translating the global remaining carbon budget to country allocations is thus not a science-driven choice, but one that represents an interplay and continuous discussion between ethics, justice, society and geophysics^{103,104} (Box 3 figure). Currently, when either implicitly or explicitly selecting a fairness principle, countries almost exclusively choose an approach that provides them with a disproportionately large share of the remaining carbon budget when seen from the perspective of another country⁷¹. This suggests an ongoing need for international cooperation and oversight to achieve consistency between national emissions budgets and international climate targets.

carbon budget is sensitive to non-CO₂ emission scenarios, it does not dictate a particular CO₂ mitigation pathway over time. This, in turn, highlights the important question of whether CO₂ emissions exceeding the budget can be reversed via carbon dioxide removal (CDR) technologies. While the global climate response to positive and negative CO₂ emissions has been shown to be approximately symmetrical for moderate amounts of negative emissions^{32,61}, CDR technologies are expensive and challenging to deploy at scale^{62,63}, and will also need to remove CO₂ that oceans and lands will release again in response to declining atmospheric CO₂^{64–66}. Furthermore, the technologies used to produce and remove CO₂ emissions will probably not produce (or remove) the same types or quantities of co-emitted non-CO₂ emissions. Current scientific understanding therefore suggests that it will be neither easy nor necessarily possible to achieve the level of CDR required to swiftly reverse the effect of substantial emissions in excess of the available budget^{66,67}, particularly with respect to long-timescale responses such as sea level⁶⁶ or other changes in the marine environment^{68–70}.

Finally, there remain substantial challenges associated with how to equitably share the remaining carbon budget among nations^{53,71}.

Issues of fairness have been interpreted differently across nations, with the result that the sum of all current national targets would produce emissions that exceed the global budget for limiting warming to 1.5 °C or “well below 2 °C”^{53,71}. This in turn suggests that many of the countries who have presented their current targets as “fair and ambitious”⁴⁴ have in fact adopted targets that are neither. To achieve a coherent set of national allocations that reflect principles of fairness and are also consistent with the global budget, it will be essential to empower and strengthen international cooperation to achieve an iterative process of evaluating and strengthening national carbon budgets.

Remaining carbon budgets are a powerful conceptual tool with clear potential to inform climate policy. Estimates of the remaining carbon budget can be used as a benchmark for international targets, and as a rationale for setting and monitoring progress towards net-zero national CO₂ emissions targets. The latest estimates of the remaining carbon budget suggest that while the global budget is small and rapidly decreasing, there is nevertheless a reasonable chance that the Paris Agreement goals remain within reach. However, this window of opportunity is closing with each passing

year of tentative and insufficient action. Halting climate change at acceptable levels will require large and rapid increases in effort on the part of both international and national players in the climate mitigation challenge.

Data availability

SR1.5 scenarios have been made available through refs. ^{87,88} at <https://data.ene.iiasa.ac.at/iamc-1.5c-explorer/>.

Code availability

The MAGICC7 model emulator is available from Z.R.J.N. upon request. Codes for producing the figures are available from H.D.M. or K.B.T. upon request.

Received: 2 October 2019; Accepted: 28 October 2020;

Published online: 30 November 2020

References

- Rogelj, J., Forster, P. M., Kriegler, E., Smith, C. J. & Séférian, R. Estimating and tracking the remaining carbon budget for stringent climate targets. *Nature* **571**, 335–342 (2019).
- Tokarska, K. B. et al. Recommended temperature metrics for carbon budget estimates, model evaluation and climate policy. *Nat. Geosci.* **12**, 964–971 (2019).
- Raupach, M. R. et al. Sharing a quota on cumulative carbon emissions. *Nat. Clim. Change* **4**, 873–879 (2014).
- Gignac, R. & Matthews, H. D. Allocating a 2 °C cumulative carbon budget to countries. *Environ. Res. Lett.* **10**, 075004 (2015).
- Nauels, A. et al. ZERO IN ON the Remaining Carbon Budget and Decadal Warming Rates. *The CONSTRAIN Project Annual Report 2019* (2019); <https://doi.org/10.5518/100/20>
- Matthews, H. D. & Caldeira, K. Stabilizing climate requires near-zero emissions. *Geophys. Res. Lett.* **35**, L04705 (2008).
- IPCC in *Special Report on Global Warming of 1.5 °C* (eds Masson-Delmotte, V. et al.) Ch. 2 (WMO, 2018).
- Mengis, N., Partanen, A.-I., Jalbert, J. & Matthews, H. D. 1.5 °C carbon budget dependent on carbon cycle uncertainty and future non-CO₂ forcing. *Sci. Rep.* **8**, 5831 (2018).
- Tokarska, K. B., Gillett, N. P., Arora, V. K., Lee, W. G. & Zickfeld, K. The influence of non-CO₂ forcings on cumulative carbon emissions budgets. *Environ. Res. Lett.* **13**, 034039 (2018).
- Matthews, H. D. et al. Estimating carbon budgets for ambitious climate targets. *Curr. Clim. Change Rep.* **3**, 69–77 (2017).
- Millar, R. J. & Friedlingstein, P. The utility of the historical record for assessing the transient climate response to cumulative emissions. *Philos. Trans. R. Soc. A* **376**, 20160449 (2018).
- Matthews, H. D., Gillett, N. P., Stott, P. A. & Zickfeld, K. The proportionality of global warming to cumulative carbon emissions. *Nature* **459**, 829–832 (2009).
- Zickfeld, K., Arora, V. K. & Gillett, N. P. Is the climate response to CO₂ emissions path dependent? *Geophys. Res. Lett.* **39**, L05703 (2012).
- Mengis, N. & Matthews, H. D. Non-CO₂ forcing changes will likely decrease the remaining carbon budget for 1.5°C. *NPL. Clim. Atmos. Sci.* **3**, 19 (2020).
- Matthews, H. D. et al. An integrated approach to quantifying uncertainties in the remaining carbon budget. *Commun. Earth Environ.* (in the press).
- Arora, V. K. et al. Carbon-concentration and carbon-climate feedbacks in CMIP6 models and their comparison to CMIP5 models. *Biogeosciences* **17**, 4173–4222 (2020).
- Jones, C. D. & Friedlingstein, P. Quantifying process-level uncertainty contributions to TCRE and carbon budgets for meeting Paris Agreement climate targets. *Environ. Res. Lett.* **15**, 074019 (2020).
- Tokarska, K. B. et al. Past warming trend constrains future warming in CMIP6 models. *Sci. Adv.* **6**, eaaz9549 (2020).
- Jiménez-de-la-Cuesta, D. & Mauritsen, T. Emergent constraints on Earth's transient and equilibrium response to doubled CO₂ from post-1970s global warming. *Nat. Geosci.* **12**, 902–905 (2019).
- Leduc, M., Matthews, H. D. & de Elía, R. Quantifying the limits of a linear temperature response to cumulative CO₂ emissions. *J. Clim.* **28**, 9955–9968 (2015).
- Tokarska, K. B., Gillett, N. P., Weaver, A. J., Arora, V. K. & Eby, M. The climate response to five trillion tonnes of carbon. *Nat. Clim. Change* **6**, 851–855 (2016).
- Leduc, M., Matthews, H. D. & de Elía, R. Regional estimates of the transient climate response to cumulative CO₂ emissions. *Nat. Clim. Change* **6**, 474–478 (2016).
- Herrington, T. & Zickfeld, K. Path independence of climate and carbon cycle response over a broad range of cumulative carbon emissions. *Earth Syst. Dynam.* **5**, 409–422 (2014).
- Winton, M., Takahashi, K. & Held, I. M. Importance of ocean heat uptake efficacy to transient climate change. *J. Clim.* **23**, 2333–2344 (2010).
- Armour, K. C., Bitz, C. M. & Roe, G. H. Time-varying climate sensitivity from regional feedbacks. *J. Clim.* **26**, 4518–4534 (2013).
- Andrews, T. et al. Accounting for changing temperature patterns increases historical estimates of climate sensitivity. *Geophys. Res. Lett.* **45**, 8490–8499 (2018).
- Gasser, T. et al. Path-dependent reductions in CO₂ emission budgets caused by permafrost carbon release. *Nat. Geosci.* **11**, 830–835 (2018).
- Comyn-Platt, E. et al. Carbon budgets for 1.5 and 2 °C targets lowered by natural wetland and permafrost feedbacks. *Nat. Geosci.* **11**, 568–573 (2018).
- Gillett, N. P., Arora, V. K., Matthews, D. & Allen, M. R. Constraining the ratio of global warming to cumulative CO₂ emissions using CMIP5 simulations. *J. Clim.* **26**, 6844–6858 (2013).
- IPCC in *Climate Change 2013: The Physical Science Basis* (eds Stocker, T. F. et al.) 1029–1136 (Cambridge Univ. Press, 2013).
- MacDougall, A. H., Zickfeld, K., Knutti, R. & Matthews, H. D. Sensitivity of carbon budgets to permafrost carbon feedbacks and non-CO₂ forcings. *Environ. Res. Lett.* **10**, 125003 (2015).
- Tokarska, K. B., Zickfeld, K. & Rogelj, J. Path independence of carbon budgets when meeting a stringent global mean temperature target after an overshoot. *Earth's Future* **7**, 1283–1295 (2019).
- MacDougall, A. H. et al. Is there warming in the pipeline? A multi-model analysis of the Zero Emissions Commitment from CO₂. *Biogeosciences* **17**, 2987–3016 (2020).
- Hienola, A. et al. The impact of aerosol emissions on the 1.5 °C pathways. *Environ. Res. Lett.* **13**, 044011 (2018).
- Lelieveld, J. et al. Effects of fossil fuel and total anthropogenic emission removal on public health and climate. *Proc. Natl Acad. Sci. USA* **116**, 7192–7197 (2019).
- Rogelj, J. et al. Air-pollution emission ranges consistent with the representative concentration pathways. *Nat. Clim. Change* **4**, 446–450 (2014).
- Rogelj, J. et al. Mitigation choices impact carbon budget size compatible with low temperature goals. *Environ. Res. Lett.* **10**, 075003 (2015).
- Rogelj, J. et al. Differences between carbon budget estimates unravelled. *Nat. Clim. Change* **6**, 245–252 (2016).
- Haustein, K. et al. A real-time global warming index. *Sci. Rep.* **7**, 15417 (2017).
- Rogelj, J., Schleussner, C.-F. & Hare, W. Getting it right matters: temperature goal interpretations in geoscience research. *Geophys. Res. Lett.* **44**, 10662–10665 (2017).
- United Nations Framework Convention on Climate Change (United Nations, 1992); <https://unfccc.int/resource/docs/convkp/conveng.pdf>
- Schleussner, C.-F. et al. Science and policy characteristics of the Paris Agreement temperature goal. *Nat. Clim. Change* **6**, 827–835 (2016).
- Tokarska, K. B. et al. Uncertainty in carbon budget estimates due to internal climate variability. *Environ. Res. Lett.* **15**, 104064 (2020).
- Adoption of the Paris Agreement FCCC/CP/2015/L.9/Rev.1 (UNFCCC, 2015).
- Knutti, R. & Rogelj, J. The legacy of our CO₂ emissions: a clash of scientific facts, politics and ethics. *Clim. Change* **133**, 361–373 (2015).
- IPCC in *Global Warming of 1.5 °C* (eds Masson-Delmotte, V. et al.) Ch. 1 (WMO, 2018).
- Hawkins, E. et al. Estimating changes in global temperature since the preindustrial period. *Bull. Am. Meteorol. Soc.* **98**, 1841–1856 (2017).
- Schurer, A. P., Mann, M. E., Hawkins, E., Tett, S. F. B. & Hegerl, G. C. Importance of the pre-industrial baseline for likelihood of exceeding Paris goals. *Nat. Clim. Change* **7**, 563–568 (2017).
- Richardson, M., Cowtan, K., Hawkins, E. & Stolpe, M. B. Reconciled climate response estimates from climate models and the energy budget of Earth. *Nat. Clim. Change* **6**, 931–935 (2016).
- The Emissions Gap Report 2019 (United Nations Environment Programme, 2019); <https://go.nature.com/3erYx1u>
- Rogelj, J. et al. Understanding the origin of Paris Agreement emission uncertainties. *Nat. Commun.* **8**, 15748 (2017).
- den Elzen, M., Janssen, M., Rotmans, J., Swart, R. & Vries, B. Allocating constrained global carbon budgets: inter-regional and inter-generational equity for a sustainable world. *Int. J. Glob. Energy Issues* **4**, 287–301 (1992).
- Robiou du Pont, Y., Jeffery, M. L., Gütschow, J., Christoff, P. & Meinshausen, M. National contributions for decarbonizing the world economy in line with the G7 agreement. *Environ. Res. Lett.* **11**, 054005 (2016).
- Höhne, N., den Elzen, M. & Escalante, D. Regional GHG reduction targets based on effort sharing: a comparison of studies. *Clim. Policy* **14**, 122–147 (2014).

55. Gibson, R. B. et al. *From Paris to Projects: Clarifying the Implications of Canada's Climate Change Mitigation Commitments for the Planning and Assessment of Projects and Strategic Undertakings* (University of Waterloo, 2019).
56. Crownshaw, T. et al. Over the horizon: exploring the conditions of a post-growth world. *Anthr. Rev.* **6**, 117–141 (2019).
57. Smith, C. J. et al. Current fossil fuel infrastructure does not yet commit us to 1.5 °C warming. *Nat. Commun.* **10**, 101 (2019).
58. Tong, D. et al. Committed emissions from existing energy infrastructure jeopardize 1.5 °C climate target. *Nature* **572**, 373–377 (2019).
59. Matthews, H. D. Quantifying historical carbon and climate debts among nations. *Nat. Clim. Change* **6**, 60–64 (2016).
60. Rogelj, J. et al. A new scenario logic for the Paris Agreement long-term temperature goal. *Nature* **573**, 357–363 (2019).
61. Zickfeld, K., MacDougall, A. H. & Matthews, H. D. On the proportionality between global temperature change and cumulative CO₂ emissions during periods of net negative CO₂ emissions. *Environ. Res. Lett.* **11**, 055006 (2016).
62. Smith, P. et al. Biophysical and economic limits to negative CO₂ emissions. *Nat. Clim. Change* **6**, 42–50 (2016).
63. Fuss, S. et al. Negative emissions—part 2: costs, potentials and side effects. *Environ. Res. Lett.* **13**, 063002 (2018).
64. Cao, L. & Caldeira, K. Atmospheric carbon dioxide removal: long-term consequences and commitment. *Environ. Res. Lett.* **5**, 024011 (2010).
65. Jones, C. D. et al. Simulating the Earth system response to negative emissions. *Environ. Res. Lett.* **11**, 095012 (2016).
66. Tokarska, K. B. & Zickfeld, K. The effectiveness of net negative carbon dioxide emissions in reversing anthropogenic climate change. *Environ. Res. Lett.* **10**, 094013 (2015).
67. Nemet, G. F. et al. Negative emissions—part 3: Innovation and upscaling. *Environ. Res. Lett.* **13**, 063003 (2018).
68. Frölicher, T. L. & Joos, F. Reversible and irreversible impacts of greenhouse gas emissions in multi-century projections with the NCAR global coupled carbon cycle-climate model. *Clim. Dyn.* **35**, 1439–1459 (2010).
69. Mathesius, S., Hofmann, M., Caldeira, K. & Schellnhuber, H. J. Long-term response of oceans to CO₂ removal from the atmosphere. *Nat. Clim. Change* **5**, 1107–1113 (2015).
70. Li, X., Zickfeld, K., Mathesius, S., Kohfeld, K. & Matthews, J. B. R. Irreversibility of marine climate change impacts under carbon dioxide removal. *Geophys. Res. Lett.* **47**, e2020GL088507 (2020).
71. Meinshausen, M. et al. National post-2020 greenhouse gas targets and diversity-aware leadership. *Nat. Clim. Change* **5**, 1098–1106 (2015).
72. Morice, C. P., Kennedy, J. J., Rayner, N. A. & Jones, P. D. Quantifying uncertainties in global and regional temperature change using an ensemble of observational estimates: the HadCRUT4 data set. *J. Geophys. Res. Atmospheres* **117**, D08101 (2012).
73. Cowtan, K. *Coverage Bias in the HadCRUT4 Temperature Series and its Impact on Recent Temperature Trends. UPDATE: COBE-SST2 Based Land-Ocean Dataset* (2017); <https://www-users.york.ac.uk/~kdc3/papers/coverage2013/update.171107.pdf>
74. Cowtan, K. et al. Robust comparison of climate models with observations using blended land air and ocean sea surface temperatures. *Geophys. Res. Lett.* **42**, 6526–6534 (2015).
75. Pfeleiderer, P., Schleussner, C.-F., Mengel, M. & Rogelj, J. Global mean temperature indicators linked to warming levels avoiding climate risks. *Environ. Res. Lett.* **13**, 064015 (2018).
76. Schurer, A. et al. Estimating the Transient Climate Response from Observed Warming. *J. Clim.* **31**, 8645–8663 (2018).
77. Meinshausen, M. et al. Greenhouse-gas emission targets for limiting global warming to 2 °C. *Nature* **458**, 1158–1162 (2009).
78. Kumar, S. et al. Land use/cover change impacts in CMIP5 climate simulations: a new methodology and 21st century challenges. *J. Geophys. Res. Atmospheres* **118**, 6337–6353 (2013).
79. Simmons, C. T. & Matthews, H. D. Assessing the implications of human land-use change for the transient climate response to cumulative carbon emissions. *Environ. Res. Lett.* **11**, 035001 (2016).
80. Lawrence, D. M. et al. The Land Use Model Intercomparison Project (LUMIP) contribution to CMIP6: rationale and experimental design. *Geosci. Model Dev.* **9**, 2973–2998 (2016).
81. IPCC in *Climate Change 2013: The Physical Science Basis* (eds T. F. Stocker et al.) 33–115 (Cambridge Univ. Press, 2013).
82. Millar, R. J. et al. Emission budgets and pathways consistent with limiting warming to 1.5 °C. *Nat. Geosci.* **10**, 741–747 (2017).
83. Tokarska, K. B. & Gillett, N. P. Cumulative carbon emissions budgets consistent with 1.5 °C global warming. *Nat. Clim. Change* **8**, 296–299 (2018).
84. Frölicher, T. L. & Paynter, D. J. Extending the relationship between global warming and cumulative carbon emissions to multi-millennial timescales. *Environ. Res. Lett.* **10**, 075002 (2015).
85. Koven, C. D., Lawrence, D. M. & Riley, W. J. Permafrost carbon–climate feedback is sensitive to deep soil carbon decomposability but not deep soil nitrogen dynamics. *Proc. Natl Acad. Sci. USA* **112**, 3752–3757 (2015).
86. McGuire, A. D. et al. Dependence of the evolution of carbon dynamics in the northern permafrost region on the trajectory of climate change. *Proc. Natl Acad. Sci. USA* **115**, 3882–3887 (2018).
87. Huppmann, D., Rogelj, J., Kriegler, E., Krey, V. & Riahi, K. A new scenario resource for integrated 1.5 °C research. *Nat. Clim. Change* **8**, 1027–1030 (2018).
88. Huppmann, D. et al. *IAMC 1.5°C Scenario Explorer and Data hosted by IIASA* (IIASA, 2018); <https://doi.org/10.22022/SR15/08-2018.15429>
89. Meinshausen, M., Raper, S. C. B. & Wigley, T. M. L. Emulating coupled atmosphere-ocean and carbon cycle models with a simpler model, MAGICC6 – Part 1: Model description and calibration. *Atmos. Chem. Phys.* **11**, 1417–1456 (2011).
90. Meinshausen, M. et al. The shared socio-economic pathway (SSP) greenhouse gas concentrations and their extensions to 2500. *Geosci. Model Dev.* **13**, 3571–3605 (2020).
91. Friedlingstein, P. et al. Global Carbon Budget 2019. *Earth Syst. Sci. Data* **11**, 1783–1838 (2019).
92. Allen, M. R. et al. Warming caused by cumulative carbon emissions towards the trillionth tonne. *Nature* **458**, 1163–1166 (2009).
93. Zickfeld, K., Eby, M., Matthews, H. D. & Weaver, A. J. Setting cumulative emissions targets to reduce the risk of dangerous climate change. *Proc. Natl Acad. Sci. USA* **106**, 16129–16134 (2009).
94. Eyring, V. et al. Overview of the Coupled Model Intercomparison Project Phase 6 (CMIP6) experimental design and organization. *Geosci. Model Dev.* **9**, 1937–1958 (2016).
95. Taylor, K. E., Stouffer, R. J. & Meehl, G. A. An overview of CMIP5 and the experiment design. *Bull. Am. Meteorol. Soc.* **93**, 485–498 (2011).
96. Forster, P. M., Maycock, A. C., McKenna, C. M. & Smith, C. Latest climate models confirm need for urgent mitigation. *Nat. Clim. Change* **10**, 7–10 (2020).
97. Sutton, R. T. ESD Ideas: a simple proposal to improve the contribution of IPCC WGI to the assessment and communication of climate change risks. *Earth Syst. Dynam.* **9**, 1155–1158 (2018).
98. IPCC in *Climate Change 2013: The Physical Science Basis. Summary for Policymakers* (eds Stocker, T. F. et al.) (Cambridge Univ. Press, 2013).
99. Jones, C. D. et al. The Zero Emissions Commitment Model Intercomparison Project (ZECMIP) contribution to C4MIP: quantifying committed climate changes following zero carbon emissions. *Geosci. Model Dev.* **12**, 4375–4385 (2019).
100. Forster, P. M. et al. Evaluating adjusted forcing and model spread for historical and future scenarios in the CMIP5 generation of climate models. *J. Geophys. Res. Atmospheres* **118**, 1139–1150 (2013).
101. Grose, M. R., Gregory, J., Colman, R. & Andrews, T. What climate sensitivity index is most useful for projections? *Geophys. Res. Lett.* **45**, 1559–1566 (2018).
102. Höhne, N., den Elzen, M. & Escalante, D. Regional GHG reduction targets based on effort sharing: a comparison of studies. *Clim. Policy* **14**, 122–147 (2014).
103. McKinnon, C. Climate justice in a carbon budget. *Clim. Change* **133**, 375–384 (2015).
104. Samson, J., Berteaux, D., McGill, B. J. & Humphries, M. M. Geographic disparities and moral hazards in the predicted impacts of climate change on human populations. *Glob. Ecol. Biogeogr.* **20**, 532–544 (2011).

Acknowledgements

We are grateful for the opportunity to have discussed these and other issues at the International Workshop on the Remaining Carbon Budget, organized with the support of the Global Carbon Project, the CRESCENDO project, Stanford University, the University of Melbourne, and Simon Fraser University. H.D.M. has been supported by funding from the Concordia University Research Chair programme and the Natural Sciences and Engineering Research Council of Canada (NSERC Discovery Grant RGPIN-2017-04159). K.B.T., J.R., P.M.F., R.K. and R.S. were supported by the European Union's Horizon 2020 research and innovation programme under grant agreement No. 820829 (CONSTRAIN project). J.G.C. was supported by the Australian National Environmental Science Program – Earth Systems and Climate Change Hub. P.F. and T.L.F. were supported by the European Union's Horizon 2020 research and innovation programme under grant agreement No. 821003 (4C project). T.L.F. was also supported by the Swiss National Science Foundation under grant PP00P2_170687. A.H.M. and K.Z. are supported by the Natural Sciences and Engineering Research Council of Canada Discovery Grant Program. C.D.J. was supported by the Joint UK BEIS/Defra Met Office Hadley Centre Climate Programme (GA01101) and by H2020 EU project CRESCENDO under grant agreement No. 641816. R.B.J. and J.G.C. acknowledge support from the Gordon and Betty Moore Foundation (GBMF5439). C.K. is supported by the US DOE, BER, RGMA program through the ECRP and RUBISCO projects.

Author contributions

H.D.M. initiated the study and wrote the manuscript with input from K.B.T., Z.R.J.N., J.R., M.M., N.M., J.G.C., T.L.F. and suggestions from other authors. H.D.M., K.B.T. and Z.R.J.N. made the figures. All authors participated in discussions at the International Workshop on the Remaining Carbon budget which initiated this work, as well as in manuscript editing and revisions.

Competing interests

The authors declare no competing interests.

Additional information

Correspondence should be addressed to H.D.M.

Peer review information Primary handling editors: Tamara Goldin; Heike Langenberg; Xujia Jiang

Reprints and permissions information is available at www.nature.com/reprints.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

© Springer Nature Limited 2020